


Transmission profile of the Dutch Open Telescope H α Lyot filter

J. Koza, R.H. Hammerschlag *, J. Rybák, P. Gömöry, A. Kučera, P. Schwartz

Astronomical Institute, Slovak Academy of Sciences, Tatranská Lomnica, Slovakia

* Leiden Observatory, Leiden, The Netherlands




SLOVAK RESEARCH
AND DEVELOPMENT
AGENCY




Imaging spectroscopy

quasi-simultaneous acquisition of wide-field images in many wavelengths in a spectral line profile

Dutch Open Telescope H α Lyot filter at $\Delta\lambda = -0.8 \quad -0.6 \quad -0.4 \quad -0.2 \quad 0 \quad 0.2 \quad 0.4 \quad 0.6 \quad 0.8 \text{ \AA}$



Imaging spectroscopy



Instruments employing:

Lyot filter

Dutch Open Telescope ($H\alpha$, Ba II 4554 Å)

Narrowband Filter Imager at Hinode

CoMP at Mauna Loa Observatory

CoMP-S at Lomnický Peak Observatory

ChroTel at Tenerife

ChroMag at High Altitude Observatory

Fabry – Pérot interferometer


IBIS at Dunn Solar Telescope

CRISP at Swedish 1-m Solar Telescope

GFPI at GREGOR

TESOS at Vacuum Tower Telescope

Lyot filter in brief





- a sequence of polarizers P and birefringent crystals of thickness $e, 2e, 4e, 8e, 16e, \dots$
- periodic transmission profile with multiple peaks
- selection of a spectral range by a broad-band prefilter

Theoretical transmission profile:

$$T = \prod_{k=1}^N \cos^2 \frac{2^{k-1} \pi e J}{\lambda} = \frac{\sin^2 \frac{2^N \pi e J}{\lambda}}{4^N \sin^2 \frac{\pi e J}{\lambda}}$$


- N – number of polarizer-crystal stages
- e – thickness of the thinnest crystal plate
- J – birefringence of crystal




In practice, the transmission profile of a single peak is often **approximated by a Gaussian**.

Application of the DOT H α Lyot filter transmission

Comparing of **observations** with results of **simulations**




DOT observation of bright points in 2004 Oct 6 in the blue wing of H α at $\Delta\lambda = -0.8 \text{ \AA}$.



Emergent intensity in the H α blue wing at $\Delta\lambda = -0.8 \text{ \AA}$ resulting from 3D MHD LTE simulation after convolving with the DOT Lyot filter transmission approximated by Gaussian (left) and Airy function (right).

Aims

Understanding an instrument may be vital for correct interpreting of data.




the DOT H α Lyot filter in deployment

[Bettonvil et al. 2006: Proc. SPIE Conf. Series 6269](#)

Aims

Understanding an instrument may be vital for correct interpreting of data.


- to summarize available facts about the transmission profile of the DOT H α Lyot filter
- to confront them with observations and suspicions on the leak of parasitic light
- to reconcile discrepancy between limb observations and spectroscopic measurement of filter
- to present a method for indirect testing of transmission profiles of Lyot filters
- two new theoretical transmission profiles of the DOT H α Lyot filter




the DOT H α Lyot filter in deployment

Problem 1: DOT limb image *versus* images from other instruments


DOT H α center
speckle-reconstructed image



FPI / VTT H α center
AO, raw image



CRISP / SST H α +0.3 Å
AO + MOMFBD



[Rutten 2007: ASPC 368, 27](#)
[Rutten 2013: ASPC 470, 49](#)

[Puschmann et al. 2006: A&A 451, 1151](#)

[Antolin & Rouppe van der Voort 2012: ApJ 745, 152](#)



“I doubt that the double limb is caused by parasitic light (continuum leak outside the H α passband).”
[from “Observing the Solar Chromosphere” \(Rutten 2007: ASPC 368, 27\)](#)

“The sharp limb is probably parasitic continuum light (cf. Bray & Loughhead 1974).”
[from “Twists to Solar Spicules” \(Rutten 2013: ASPC 470, 49\)](#)

**DOT limb image shows the sharp limb what is not the case of images from the other instruments.
The DOT sharp limb may be due to parasitic light.**

Problem 2: DOT limb image *versus* spectroscopic measurement of filter

DOT H α center
speckle-reconstructed image



Measured transmission profile of the DOT H α Lyot filter.



The spectroscopic measurement of the filter showed:

- almost symmetric and Gaussian-like transmission profile with FWHM = 250 mÅ without subsidiary maxima or far-center sidelobes **ruling out a leak of unwanted parasitic light**
- invariance of the profile in tuning

[Rutten 2007: ASPC 368, 27](#)

[Rutten 2013: ASPC 470, 49](#)

Indirect testing the filter transmission



- Large discrepancy of observed and anticipated ratios.
- An indication of presence of parasitic light in DOT H α images supporting the suspicions in Rutten 2007 and 2013?


DOT observation of quiet Sun
at disk center 2005 Oct 19
taken by Hitachi camera

similar scene on 2007 Sep 28
but taken by RedLake camera


Observed *versus* anticipated wing-to-center ratios

Ratio	DOT H α Observations		Atlas H α Profile + + Gaussian
	2005 Oct 19	2007 Sep 28	
$E(\pm 0.7) / E(0)$	2.32	2.34	3.28
$E(\pm 0.7) / E(\pm 0.35)$	1.75	1.75	2.10
$E(\pm 0.35) / E(0)$	1.33	1.34	1.56

Indirect testing the filter transmission



approximation for transmission profile of a single peak of Lyot filter



$$T = \text{sinc}^2(\Delta\lambda) = \left(\frac{\sin \pi x}{\pi x} \right)^2, x = 2k \frac{\Delta\lambda}{FWHM}$$



[Gaizauskas 1976, JRASC 70, 1](#)
[Koza et al. 2014: AN 335, 409](#)

- Gaussian and sinc² models yield ratios significantly exceeding the observed ones.
- It suggests in contradiction with the spectroscopic measurement of the filter that its real transmission profile might have larger throughput than these models.

Observed *versus* anticipated wing-to-center ratios

Ratio	DOT Hα Observations		Atlas Hα Profile +	
	2005 Oct 19	2007 Sep 28	+ Gaussian	+ sinc ²
$E(\pm 0.7) / E(0)$	2.32	2.34	3.28	2.78
$E(\pm 0.7) / E(\pm 0.35)$	1.75	1.75	2.10	1.94
$E(\pm 0.35) / E(0)$	1.33	1.34	1.56	1.43

New transmission models



Ad hoc rectangular extensions Λ and Π of the Gaussian and sinc^2 function at $\Delta\lambda = \pm 2 \text{ \AA}$

Rectangle	Area [mÅ]	Width [mÅ]	Height
Λ	20.0	141	0.141
Π	11.5	107	0.107


Observed *versus* anticipated wing-to-center ratios

Ratio	DOT H α Observations		Atlas H α Profile +			
	2005 Oct 19	2007 Sep 28	+ Gaussian	+ sinc^2	+ Gaussian + Λ	+ sinc^2 + Π
$E(\pm 0.7) / E(0)$	2.32	2.34	3.28	2.78	2.35	2.35
$E(\pm 0.7) / E(\pm 0.35)$	1.75	1.75	2.10	1.94	1.77	1.76
$E(\pm 0.35) / E(0)$	1.33	1.34	1.56	1.43	1.33	1.34


Discussion

Discriminating a source of the parasitic light

Transmission of prefilter (FWHM = 14.9 Å)
of the DOT H α Lyot filter



The logarithm of the prefilter transmission
(The transmission decreases upward)



Symptoms of a leak of the parasitic light:

1. dissimilarity of limb images
2. discrepancies of the wing-to-center ratios

Two likely gateways:

1. **IR window** in the prefilter transmission (broadband transmission of $10^{-5.5}$ at 870 - 1200 nm)
2. the main passband of the Lyot filter itself

Considering an extreme situation that the polarisers in the Lyot filter are completely ineffective in IR wavelengths, estimated IR leak ϵ_{IR} is only 1.3% from the total parasitic light.

$$\epsilon_{IR} \approx \frac{F_{IR}}{F_{H\alpha}} \frac{P_{IR}}{P_{H\alpha}} \frac{S_{IR}}{S_{H\alpha}} \Delta\lambda_{IR}$$

Then virtually all parasitic light leaks through the main passband of the prefilter and the Lyot filter.

